I'm human

Gel electrophoresis simulation worksheet answers

Gel electrophoresis basics worksheet answer key. Gel electrophoresis simulation questions. Gel electrophoresis worksheet. Gel electrophoresis simulation worksheet. Gel electrophoresis worksheet answer key. Gel electrophoresis practice worksheet answers.

As you were browsing, our system detected that something might be off. It's possible that you're a power user moving too fast, or maybe you've disabled cookies or JavaScript is blocked by a third-party plugin. Check out this support article for more info. To get back in, make sure to enable cookies and JavaScript before reloading the page. MODULE 15: GEL ELECTROPHORESIS Instructions: Complete lab exercises based on lecture material, level 3 simulation. Submit completed worksheet (20 points). 1. Ladders are used in agarose and SDS-PAGE gels. Describe purposes and compositions: Agarose gel: A diagnostic tool for visualizing fragments, separating them by size. SDS-PAGE gel: Separates proteins based on length and molecular weight, an inexpensive and accurate method. 2. In SDS-PAGE, the following chemicals are added to the protein sample before loading. Describe their purposes: Sodium dodecyl sulfate: Sets conditions for electrophoresis, ensuring all proteins separate into same-sized bands. Beta-mercaptoethanol: A reducing agent that breaks disulfide bridges. Bromophenol blue: A tracking dye that moves with the protein sample, marking its leading edge. Glycerol: Increases sample density to ensure wells sink during loading. 3. In SDS-PAGE, the separating gel is pH 8.8 and the stacking gel is pH 6.8. Explain why they're different and how they work: Stacking gel: Lines up protein samples before entering the resolving gel at the same time. Separating gel: Separates samples based on molecular weight, allowing for distinct bands. 4. Sodium dodecyl sulfate is present in both the gel and running buffer. Why? Ans: To make proteins negatively charged, denature them, and transport to the electrophoretic field at a rate equal to their length. In a lab analysis, agarose and SDS-PAGE gels are used to separate molecules based on size and charge. For specific types of molecules, one gel type is preferred over the other. - Plasmid: Agarose - cDNA (complementary DNA): Agarose - mRNA (messenger RNA): Agarose - Enzyme: SDS-PAGE - Kinase: SDS-PAGE